

## Geometry

Jonathan L.F. King  
 University of Florida, Gainesville FL 32611-2082, USA  
 squash@ufl.edu  
 Webpage <http://www.math.ufl.edu/~squash/>  
 23 March, 2012 (at 09:26)

ABSTRACT: Heron's theorem on the area of a triangle.  
 Maximum area of an articulated polygon.  
 Edgar's way to get an equation of a circle.

### The Triangle

Let  $\mathbf{T}$  be the triangle  $\triangle ABC$ . Following the usual convention,  $\angle A$  or just  $A$  itself will also denote the interior angle at vertex  $A$ . The edge opposite vertex  $A$  is lowercase  $\mathbf{a}$ , etc. Also,  $\mathbf{a}$  denotes the *length* of edge  $\mathbf{a}$ .

**Tools.** Recall the Law of Cosines, which asserts that

$$\text{LoCos:} \quad c^2 = a^2 + b^2 - [2ab \cdot \cos(C)].$$

Also note that

$$1: \quad \text{Area}(\triangle ABC) = \frac{1}{2}ab \cdot \sin(C),$$

since  $b \cdot \sin(C)$  is  $\text{Len}(A\text{-altitude})$ ; ie., down to edge  $\mathbf{a}$ .

**2: Heron's formula.** Fix  $\mathbf{T} := \triangle ABC$ . Then

$$2i: \quad [4 \cdot \text{Area}(\mathbf{T})]^2 = [a+b+c][-a+b+c][a-b+c][a+b-c] \\ \stackrel{\text{note}}{=} 2[a^2b^2 + b^2c^2 + a^2c^2] - [a^4 + b^4 + c^4].$$

*Equivalently (and classically),*

$$2ii: \quad \text{Area}(\mathbf{T}) = \sqrt{\sigma \cdot [\sigma - a][\sigma - b][\sigma - c]},$$

where  $\sigma := \frac{a+b+c}{2}$  is the *semi-perimeter* of  $\mathbf{T}$ .  $\diamond$

**Pf.** The Pythag thm, in form  $\sin^2 = 1^2 - \cos^2$ , gives

$$[4 \cdot \text{Area}]^2 \stackrel{\text{by (1)}}{=} [2ab \cdot \sin(C)]^2 \\ \stackrel{\text{Pythag}}{=} [2ab]^2 - [2ab \cdot \cos(C)]^2 \\ \stackrel{\text{LoCos}}{=} [2ab]^2 - [a^2 + b^2 - c^2]^2.$$

This last doesn't look symmetric in  $\mathbf{a}, \mathbf{b}, \mathbf{c}$ , but squaring, then adding, produces  $\text{RhS}(2i)$ , as desired.  $\diamond$

**Inscribed radius and circum-radius.** Let  $r$  and  $\mathcal{O}$  denote the radius and center of the in-circle of  $\mathbf{T} := \triangle ABC$ . Evidently  $\text{Area}(\triangle A\mathcal{O}B) = \frac{1}{2}rc$ . Adding this to the areas of  $\triangle B\mathcal{O}C$  and  $\triangle C\mathcal{O}A$  yields that  $\text{Area}(\mathbf{T}) = \frac{1}{2}r \cdot [a + b + c]$ . Equivalently

$$3.1: \quad \text{InRadius}(\mathbf{T}) = \frac{2 \cdot \text{Area}(\mathbf{T})}{\text{Perimeter}(\mathbf{T})}.$$

Now let  $R$  denote the radius of  $\text{CircumCircle}(\mathbf{T})$ . Let  $2\gamma$  denote the central angle  $\angle A\mathcal{O}B$  of the circle-arc not owning  $C$ . Dropping a perpendicular from  $\mathcal{O}$  to chord  $\overline{AB}$  we see that

$$\sin(\gamma) = \frac{\frac{1}{2}c}{R} = \frac{c}{2R}.$$

Thus

$$R = \frac{c}{2 \sin(\gamma)} = \frac{a \cdot b \cdot c}{2ab \cdot \sin(\gamma)}.$$

The Central-angle thm asserts that the inscribed  $\angle C$  equals  $\gamma$ . So (1) hands us

$$3.2: \quad \text{CircumRadius}(\mathbf{T}) = \frac{a \cdot b \cdot c}{4 \cdot \text{Area}(\mathbf{T})}.$$

We get this curious corollary for the radii-ratio:

$$3.3: \quad \frac{\text{CircumRadius}}{\text{InRadius}} = \frac{[a \cdot b \cdot c][a + b + c]}{8 \cdot [\text{Area}^2]}.$$

Reciprocating, then using Heron's formula (2i), gives

$$3.4: \quad \frac{2 \cdot \text{InRadius}}{\text{CircumRad}} = \frac{[-a+b+c][a-b+c][a+b-c]}{[a \cdot b \cdot c]} \\ = \left[ \frac{b+c}{a} - 1 \right] \left[ \frac{a+c}{b} - 1 \right] \left[ \frac{a+b}{c} - 1 \right].$$

### Boomerangs cannot tile a convex polygon

(Problem from David Gale.) A *boomerang* is a non-convex quadrilateral; call its  $> \frac{\pi}{2}$  interior-angle “fat”. Conversely, a quadrilat(eral) with each angle  $\leq \frac{\pi}{2}$  is a *kite*. A dissection of a polygon  $\mathbf{P}$  into *finitely many* quadrilats is a “*quadrtiling* of  $\mathbf{P}$ ”. The tiles in a quadrtiling *need not* be congruent to each other.

**4.1: Boom-Kite Thm.** *Each quadrtiling of a convex polygon  $\mathbf{P}$  must use a kite.*  $\diamond$

**4.2: Fails with “Quad” replaced by “Penta”.** Let  $\mathbf{P}$  be the square with vertices  $(\pm 2, \pm 2)$ . Cut  $\mathbf{P}$  with a polygonal path going from/to

$$(2, 2) \rightarrow (-1, 1) \rightarrow (1, -1) \rightarrow (-2, -2).$$

This cuts  $\mathbf{P}$  (which is convex) into two non-convex pentagons (which are congruent to each other).

**Exercise:** Each polygon  $\mathbf{Q}$ , convex or not, admits a (finite) tiling by non-convex *pentagons*.  $\square$

### Nested convex curves

Attributed to Archimedes is the following theorem:

**5: Theorem (Archimedes).** *Suppose  $E$  and  $D$  are compact convex sets in the plane, with  $E \supset D$ . Then the arclength (of the boundary of)  $E$  dominates the arclength of  $D$ .*  $\diamond$

**Proof.** Let a *chord* denote a line-segment having both its endpoints on  $\partial E$  and which is tangent to  $D$ . Cut off the “outside piece” of a chord from  $E$  to get the smaller convex body  $E_1 \supset D$ . Automatically,  $\text{Len}(\partial E_1)$  is less-equal that of  $\text{Len}(\partial E)$ .

We can do a sequence of cuts to get a sequence of convex bodies  $E \supset E_1 \supset E_2 \supset \dots$ , all of which are supersets of  $D$ . Moreover, we can arrange that the  $E_n$  “converge” to  $D$  –say, in the Hausdorff metric, or, even easier, in the sense that  $\bigcap_{n=1}^{\infty} E_n = D$ .

One can then show, since the objects are convex, that the arclength of  $\partial E_n$  is converging to  $\text{Len}(\partial D)$ . And  $n \mapsto \text{Len}(\partial E_n)$  is a non-increasing function.  $\diamond$

### Finding an equation of a circle, given a non-colinear triple of points.

[jk: The 4×4 Det is from edgar@mps.ohio-state.edu]

We have three points  $(\mathbf{A}, \alpha)$ ,  $(\mathbf{B}, \beta)$  and  $(\mathbf{C}, \gamma)$  in the plane and we let

$$\mathbf{E} := \begin{bmatrix} \mathbf{A} & \alpha & 1 \\ \mathbf{B} & \beta & 1 \\ \mathbf{C} & \gamma & 1 \end{bmatrix}.$$

[More generally, the coordinates can come from an arbitrary commutative ring.]

**6.1: Colinearity lemma.** *The triple of points is colinear IFF  $\text{Det}(\mathbf{E}) = 0$ .*  $\diamond$

**Pf of ( $\Rightarrow$ ).** The triple lies in a line, so the three rows of  $\mathbf{E}$ , viewed as points in  $\mathbb{R}^3$ , lie in a lift of that line to the  $z=1$  plane, hence lie in a line in  $\mathbb{R}^3$ . Consequently,  $\text{Spn}(\text{Lifted points})$  is at-most 2-dimensional.  $\diamond$

**Pf of ( $\Leftarrow$ ).** By hyp., the triple of  $\mathbf{E}$ -rows [viewed as points in  $\mathbb{R}^3$ ] lie in plane,  $\mathbf{P}$ , through the origin. But they also lie in the  $z=1$  plane; it misses the origin, so does *not equal*  $\mathbf{P}$ . Hence the intersection of these two planes lies in a line in the  $z=1$  plane; and this line projects to a line in  $\mathbb{R}^2$ .  $\diamond$

**6.2: Circle-eqn lemma.** *An equation  $\text{Func}(x, y) = 0$  of the circle through the non-colinear triple of points can be given as a 4×4 determinant-eqn:*

$$6.3: \quad \text{Det} \begin{bmatrix} x^2 + y^2 & x & y & 1 \\ \mathbf{A}^2 + \alpha^2 & \mathbf{A} & \alpha & 1 \\ \mathbf{B}^2 + \beta^2 & \mathbf{B} & \beta & 1 \\ \mathbf{C}^2 + \gamma^2 & \mathbf{C} & \gamma & 1 \end{bmatrix} = 0. \quad \diamond$$

**Proof.** Expanding LhS(6.3) along the first row shows it to be a polynomial with a common coeff for  $x^2$  and for  $y^2$  of  $\text{Det}(\mathbf{E})$ ; this latter is non-zero, courtesy (6.1). Hence (6.3) is the equation of *some* circle [which possibly is degenerate or empty].

Certainly  $(x, y) := (\mathbf{A}, \alpha)$  satisfies (6.3), since a matrix with two rows equal has  $\text{Det}=0$ . Ditto  $(\mathbf{B}, \beta)$  and  $(\mathbf{C}, \gamma)$  lie on the circle. Hence the circle is not degenerate, since a non-colinear triple lies on it.  $\diamond$

Filename: Problems/Geometry/geometry+.latex  
As of: Sunday 11Mar2012. Typeset: 23Mar2012 at 09:26.