

Inclusion-Exclusion principle

Jonathan L.F. King
 University of Florida, Gainesville FL 32611-2082, USA
 squash@ufl.edu
 Webpage <http://squash.1gainesville.com/>
 6 November, 2017 (at 00:14)

Prolegomenon. Consider a finite set, Ω , of *tokens*. and a finite indexing set \mathcal{N} ; let $N := \#\mathcal{N}$. For each $j \in \mathcal{N}$ we have a *patch*, a subset $A_j \subset \Omega$, and its complement $V_j := \Omega \setminus A_j$. Our goal is an expression for the cardinality of the union

$$\mathbf{U} := \bigcup_{j \in \mathcal{N}} A_j.$$

[N.B: This note uses $\#$ to indicate the cardinality of an index-set, and $|\cdot|$ for the cardinality of a token set.]

Notation. For each *index-set* $I \subset \mathcal{N}$, define the *patch-intersection*

$$1.1: \quad \mathbf{A}_I := \bigcap_{j \in I} A_j,$$

[and $\mathbf{V}_I := \bigcap_{j \in I} V_j$] and note^{♥1} \mathbf{A}_\emptyset is all of Ω .

Finally, let \mathcal{C}_k comprise those index-sets $I \subset \mathcal{N}$ with $\#I = k$; this, for $k = 0, 1, \dots, N$. Easily

$$1.2: \quad |\mathcal{C}_k| = \binom{N}{k}.$$

2: Inclusion-Exclusion Lemma. *With notation from above:*

$$2a: \quad |\Omega \setminus \mathbf{U}| = \sum_{k=0}^N \left[(-1)^k \cdot \sum_{I: I \in \mathcal{C}_k} |\mathbf{A}_I| \right].$$

Alternatively,

$$2b: \quad |\mathbf{U}| = \sum_{k=1}^N \left[(-1)^{k-1} \cdot \sum_{I: I \in \mathcal{C}_k} |\mathbf{A}_I| \right]. \quad \blacklozenge$$

Pf (2a) \Rightarrow (2b). Adding the RhSes causes cancellation, with only the $k=0$ term remaining. So

$$|\Omega \setminus \mathbf{U}| + \text{RhS}(2b) = (-1)^0 \cdot |\mathbf{A}_\emptyset| \stackrel{\text{note}}{=} |\Omega|.$$

So RhS(2b) equals $|\mathbf{U}|$. \blacklozenge

^{♥1}For index-sets $I, J \subset \mathcal{N}$, by definition $\mathbf{A}_I \cap \mathbf{A}_J = \mathbf{A}_{I \cup J}$. In particular, $\mathbf{A}_I \cap \mathbf{A}_\emptyset$ equals \mathbf{A}_I . I.e., \mathbf{A}_\emptyset is the identity element for intersection, on the powerset of Ω . So \mathbf{A}_\emptyset is Ω .

To establish (2a), let's prove something a little stronger. Each subset $S \subset \Omega$ yields a function $\mathbf{1}_S: \Omega \rightarrow \{0, 1\}$, the "*indicator function* of S ",

$$\mathbf{1}_S(x) := \left\{ \begin{array}{ll} 1 & \text{when } x \in S \\ 0 & \text{when } x \in \Omega \setminus S \end{array} \right\}.$$

Indicator fncs allow us to restate intersection ITOF multiplication: For a set I of indicies,

$$1.1': \quad \mathbf{1}_{\mathbf{A}_I} = \prod_{j \in I} \mathbf{1}_{A_j}.$$

Let's strengthen (2a) to equality of *functions*,

$$3.1: \quad \mathbf{1}_{\Omega \setminus \mathbf{U}} = \sum_{k=0}^N \left[(-1)^k \cdot \sum_{I: I \in \mathcal{C}_k} \mathbf{1}_{\mathbf{A}_I} \right].$$

Proof of (3.1). The RhS(3.1) equals

$$*: \quad \prod_{j \in \mathcal{N}} [1 - \mathbf{1}_{A_j}] \stackrel{\text{note}}{=} \prod_{j \in \mathcal{N}} \mathbf{1}_{V_j}.$$

After all, the LhS(*) product expands to the sum, over all subsets $I \subset \mathcal{N}$, of $\prod_{j \in I} [-\mathbf{1}_{A_j}]$. This latter product, letting $k := \#I$, equals $(-1)^k \cdot \mathbf{1}_{\mathbf{A}_I}$, courtesy (1.1').

By De Morgan's law, $\Omega \setminus \mathbf{U} = \bigcap_{j \in \mathcal{N}} V_j$. So LhS(3.1) is $\mathbf{1}_{\mathbf{V}_\mathcal{N}}$. Hence (3.1) boils down to the triviality that

$$\mathbf{1}_{\mathbf{V}_\mathcal{N}} = \prod_{j \in \mathcal{N}} \mathbf{1}_{V_j}. \quad \blacklozenge$$

4: Rem. We can rewrite (2a) in simpler [but often less convenient] form, ITOF summing over all subsets of \mathcal{N} :

$$4a: \quad |\Omega \setminus \mathbf{U}| = \sum_{I: I \subset \mathcal{N}} [(-1)^{\#I} \cdot |\mathbf{A}_I|]. \quad \square$$

5: Cardinality independence. In some combinatorial applications, cardinality $|\mathbf{A}_I|$ depends only on the number of patches being intersected. In that instance, let $F(k)$ be $\#I$, for each and every index-set I satisfying $\#I = k$. So rewrite (2a) as

$$5a: \quad \begin{aligned} |\Omega \setminus \mathbf{U}| &= \sum_{k=0}^N \left[(-1)^k \cdot \binom{N}{k} \cdot F(k) \right] \\ &= \sum_{k=0}^{\infty} \left[(-1)^k \cdot \binom{N}{k} \cdot F(k) \right], \end{aligned}$$

where, for $k > N$, our $\binom{N}{k} := \frac{[N \downarrow k]}{k!}$ is zero. \square

Incl-Excl Examples

TwoStirling numbers. Recall $\mathcal{S}(L, N)$ counts the number of partitions of an L -set into N nv-atoms.

6: TwoStirling Theorem. For all natnums L, N :

$$*: \quad \mathcal{S}(L, N) = \sum_{k=0}^N (-1)^k \cdot \frac{[N - k]^L}{k! \cdot [N - k]!} \quad \diamond$$

Proof. Let \mathcal{L} be some L -set, and \mathcal{N} an N -set; the **token** set. Previous work showed that

$$N! \cdot \mathcal{S}(L, N)$$

equals be the number of surjections $\mathcal{L} \rightarrow \mathcal{N}$. This equals the number of all fncs, minus those that miss a specified token, plus those that miss a specified pair of tokens, minus... For a subset $I \subset \mathcal{N}$, define

$$\mathbf{A}_I := \left\{ h: \mathcal{L} \rightarrow \mathcal{N} \mid \text{Range}(h) \cap I \right\}.$$

Notice that $0 \leq k \leq N$, where $k := \#I$. And that cardinality $|\mathbf{A}_I| = [N - k]^L$ depends on k , independent of the contents of I . Hence, let $F(k) := [N - k]^L$. Our (5a) reads as

$$N! \cdot \mathcal{S}(L, N) = \sum_{k=0}^N \left[(-1)^k \cdot \binom{N}{k} \cdot [N - k]^L \right].$$

Dividing by $N!$ produces (*). ♦

7: Probability of getting your own hat. The N guests leaving your party grab their hats at random from your dark closet. What does $\mathcal{P}(N)$, the probability that no one gets his own hat, tend to as $N \nearrow \infty$? ♦

Hat soln. For each subset I of your guests, let H_I denote the probability that each person in I took his own hat. It equals $H_k := [N - k]!/N!$ where k is the cardinality of I , i.e $k := \#I$. By the principle of inclusion/exclusion, our $\mathcal{P}(N)$ equals

$$\begin{aligned} H_{\emptyset} - \sum_{I: \#I=1} H_I + \sum_{I: \#I=2} H_I - \sum_{I: \#I=3} H_I + \dots + (-1)^N \sum_{I: \#I=N} H_I \\ = 1 - \binom{N}{1} H_1 + \binom{N}{2} H_2 - \binom{N}{3} H_3 + \dots + (-1)^N \binom{N}{N} H_N \\ = \sum_{k=0}^N (-1)^k \cdot \binom{N}{k} \cdot \frac{[N - k]!}{N!} = \sum_{k=0}^N \frac{(-1)^k}{k!}. \end{aligned}$$

This last is the first $k+1$ terms of the Taylor series for e^{-1} . Thus $\lim_{n \rightarrow \infty} \mathcal{P}(n)$ equals $1/e$. ♦

Derangements. A **derangement** of an N -set is a fixed-point free permutation of that set; let \mathbb{D}_N be the set of derangements. Thus

$$7a: \quad |\mathbb{D}_N| = N! \cdot \sum_{k=0}^N \frac{(-1)^k}{k!}$$

is a restatement of the above probability. □

8: Comps avoiding a part. What is $f(N)$, the number of 3-compositions of N with no PartSize=2? ♦

Remark. Bona's text asks for $f(12)$. □

Proof. Not yet typed. ♦