

## Table of Laplace transforms

Jonathan L.F. King  
squash@ufl.edu

31 March, 2018 (at 14:19)

**General Lap xforms.** For complex numbers  $\alpha$  and  $\beta$ , let  $\alpha \succ \beta$  mean  $\text{Re}(\alpha) > \text{Re}(\beta)$

With  $\mu$  an arbitrary real number and  $f, g \in \text{Ord}(\mu)$ ,<sup>♥1</sup> each formula in the table below holds for all complex  $s \succ \mu$ , unless a trailing note says otherwise.

Natural numbers:  $N, j, k$ .  
Real numbers:  $F$ . Positive reals: 7, 9.  
Complex Numbers:  $R, Q, B_j$ .

| $h$ or $h(t)$  | $\widehat{h}$ or $\widehat{h}(s)$   |
|--|---|
| <b>C:</b> $f \otimes g$  | $\widehat{f} \cdot \widehat{g}$   |
| <b>Mul:</b> $f(9t)$  | $\frac{1}{9} \cdot \widehat{f}(s/9)$                                      |
| <b>1:</b> $f'$   | $s \cdot \widehat{f}(s) - f(0)$   |
| <b>1<sub>N</sub>:</b> $f^{(N)}$  | $s^N \widehat{f}(s) - \left[ \sum_{j+k=N-1} s^j \cdot f^{(k)}(0) \right]$ |
| <b>1**<sub>Ply</sub>:</b> $[q(\mathbf{D})](f)$                                       | $q \cdot \widehat{f}$   |
| <b>1<sub>-1</sub>:</b> $\int_0^t f(x) dx \stackrel{\text{note}}{=} [1 \otimes f](t)$ | $\widehat{f}(s)/s$  |
| <b>2:</b> $f(t) \cdot t$   | $-[\widehat{f}]'$   |
| <b>2<sub>N</sub>:</b> $f(t) \cdot t^N$   | $[-1]^N \widehat{f}^{(N)}$  |
| <b>2<sub>-1</sub>:</b> $f(t)/t$  | $\int_s^\infty \widehat{f}$   |
| <b>3:</b> $f(t) \cdot e^{Rt}$  | $\widehat{f}(s - R)$  |
| <b>4:</b> $f(t-7) \cdot \mathbf{H}(t-7)$   | $e^{-7s} \cdot \widehat{f}(s)$  |
| $f(t) \cdot \mathbf{H}(t-7)$   | $e^{-7s} \cdot \widehat{f(\widehat{t+7})}(s)$                             |
| <b>5:</b> $f$ has period 7   | $\mathcal{L}_7(f)(s) / [1 - e^{-7s}]$                                     |

The “**D**” in formula (1<sub>Ply</sub>) is the differentiation operator. The “**q**” is a general degree- $N$  polynomial

$$q(s) = B_0 + B_1s + B_2s^2 + \dots + B_Ns^N.$$

This\*\* needs that derivatives  $f^{(k)}(0)$  are zero, for each  $k \in [0..N)$ . In that case, (1<sub>Ply</sub>) is says that function

$$s \mapsto q(s) \cdot \widehat{f}(s)$$

<sup>♥1</sup>See “exponential order” in the `laplace.xform.latex` file.

is the xform of  $[B_0f + B_1f' + B_2f'' + \dots + B_Nf^{(N)}]$ .  
Line (5) arranges for periodic input fncs with

$$\mathcal{L}_7(f)(s) := \int_0^7 e^{-st} f(t) dt.$$

Formula (5) holds  $\forall s \succ 0$ .

Line (3) is valid for all  $s \succ R + \mu$ .

Formula (Mul) is valid for all  $s \succ 9\mu$ .

**Specific Lap xforms.** Many of the below come from the “General” table, with  $f := \mathbf{1}$ .

| $h$ or $h(t)$                      | $\widehat{h}$ or $\widehat{h}(s)$  |
|------------------------------------|------------------------------------|
| <b>1</b>                           | $1/s$                              |
| $t^N$                              | $N! / s^{N+1}$                     |
| <b>12:</b> $t^R$                   | $\Gamma(R+1) / s^{R+1}$            |
| $1/\sqrt{t}$                       | $\sqrt{\pi} / \sqrt{s}$            |
| $\sqrt{t}$                         | $\frac{1}{2} \sqrt{\pi} / s^{3/2}$ |
| <b>13:</b> $e^{Rt} \cdot t^N$      | $N! / [s - R]^{N+1}$               |
| <b>14:</b> $e^{Rt} \cdot \sin(Ft)$ | $\frac{F}{[s-R]^2 + F^2}$          |
| $e^{Rt} \cdot \cos(Ft)$            | $\frac{s-R}{[s-R]^2 + F^2}$        |

Formula (12) requires  $R \succ -1$ , as does this defn:

$$\Gamma(R+1) := \int_0^\infty t^R e^{-t} dt.$$

The (12) formulas hold for all  $s \succ 0$ .

Formulas (13,14) are valid for all  $s \succ R$ .

Filename: `Problems/Analysis/Calculus/lapxform.table.latex`  
As of: `Wednesday 15Mar2006`. Typeset: `31Mar2018 at 14:19`.