

## Linear Recurrence using matrices

Jonathan L.F. King  
 University of Florida, Gainesville FL 32611  
 squash@ufl.edu  
 Webpage <http://squash.1gainesville.com/>  
 3 November, 2015 (at 17:18)

See also [Problems/NumberTheory/fibonacci.latex](#)

**Notation.** Use “ $\overset{r}{\sim}$ ” for row-equivalence: I.e,  $K \times N$  matrices  $M \overset{r}{\sim} M'$  IFF we can get from  $M$  to  $M'$  using row operations. Use “ $\overset{c}{\sim}$ ” for column-equivalence.

Two  $N \times N$  matrices  $B$  &  $C$  are *similar*, or *conjugate* to each other, if  $\boxed{\text{there exists}}$  an invertible matrix  $U$  such that

$$U^{-1} \cdot B \cdot U = C.$$

Use  $B \overset{\text{sim}}{\asymp} C$  for the similarity equiv-relation.

Call a matrix OTForm  $s\mathbf{I} \overset{\text{note}}{\equiv} \begin{bmatrix} s & 0 \\ 0 & s \end{bmatrix}$ , a *dilation*; its action on the plane is simply to scale-uniformly by factor  $s$ . *Every* matrix commutes with  $\mathbf{I}$ , and so:

1: A dilation is only conjugate to itself.

I.e, for invertible  $U$ , necessarily  $U^{-1} \cdot s\mathbf{I} \cdot U = s\mathbf{I}$ .

For  $T$  a square matrix (or a trn from a vectorspace to itself) and a complex number  $\alpha$ , define the subspace

2:  $\mathbb{E}_{T,\alpha} := \{\text{vectors } \mathbf{v} \mid T\mathbf{v} = \alpha\mathbf{v}\}$ .

Saying that “ $\alpha$  is a  $T$ -eigenvalue” is the same as saying that  $\text{Dim}(\mathbb{E}_{T,\alpha}) \geq 1$ .

### Linear recurrence

A fibonacci-like sequence  $\vec{z} := (z_n)_{n=-\infty}^{\infty}$  is specified using (complex) numbers  $\mathcal{S}$  and  $\mathcal{P}$ , with  $\mathcal{P}$  non-zero<sup>♥1</sup>, by

3:  $z_{n+2} := \mathcal{S}z_{n+1} - \mathcal{P}z_n,$

and some initial condition  $(z_1, z_0)$ . With

$$G := \begin{bmatrix} \mathcal{S} & -\mathcal{P} \\ 1 & 0 \end{bmatrix}, \quad \text{then } \begin{bmatrix} z_{n+1} \\ z_n \end{bmatrix} = G^n \cdot \begin{bmatrix} z_1 \\ z_0 \end{bmatrix},$$

for each integer  $n$ . We want to diagonalize  $G$ .

<sup>♥1</sup> Actually, recurrence (3) can be run backwards, giving  $z_n$  values when  $n$  is negative, as soon as  $(\mathcal{S}, \mathcal{P}) \neq (0, 0)$ . However, the  $\mathcal{P} \neq 0$  case is the interesting case.

Its char-poly is  $f(x) = f_G(x) := x^2 - \mathcal{S}x + \mathcal{P}$ . Factor this  $f$  as  $f(x) = [x - \alpha][x - \beta]$ , with  $\alpha, \beta \in \mathbb{C}$ . Equating coeffs in the polynomial gives

$$\begin{aligned} 4: \quad \alpha + \beta &= \mathcal{S}; & (\text{Sum}) \\ \alpha \cdot \beta &= \mathcal{P}. & (\text{Product}) \end{aligned}$$

Since  $\mathcal{P} \neq 0$ , necessarily  $\boxed{\alpha \neq 0}$  and  $\boxed{\beta \neq 0}$ . Each of  $\alpha, \beta$  is a root of  $f$ , hence

$$5: \quad \alpha^2 = \mathcal{S}\alpha - \mathcal{P} \quad \text{and} \quad \beta^2 = \mathcal{S}\beta - \mathcal{P}.$$

**Finding an  $\alpha$ -eigenvector.** An  $\alpha$ -evec (for  $G$ ) is an element of  $\text{Nul}(G - \alpha\mathbf{I})$ . Applying row operations,

$$6: \quad G - \alpha\mathbf{I} = \begin{bmatrix} \mathcal{S} - \alpha & -\mathcal{P} \\ 1 & -\alpha \end{bmatrix} \overset{r}{\sim} \begin{bmatrix} 1 & -\alpha \\ \mathcal{S} - \alpha & -\mathcal{P} \end{bmatrix} \overset{r}{\sim} \begin{bmatrix} 1 & -\alpha \\ 0 & 0 \end{bmatrix}.$$

This last  $\overset{r}{\sim}$  requires no computation, since the rows *must* be linearly-dependent (since  $\alpha$  is an eigenvalue of  $G$ ).

This last matrix has one free column, and evidently multiplies  $\begin{bmatrix} \alpha \\ 1 \end{bmatrix}$  to the zero-vector. So

7: The singleton  $\left\{ \begin{bmatrix} \alpha \\ 1 \end{bmatrix} \right\}$  is a basis for the  $\alpha$ -eigenspace of  $G$ , which is one-dimensional.

Instead of row-ops to show that  $\mathbb{E}_{G,\alpha}$  (the  $\alpha$ -eigenspace of  $G$ ) is one-dim'al, we could have argued as follows: Were  $\mathbb{E}_{G,\alpha}$  two-dim'al, then  $G \overset{\text{sim}}{\asymp} \begin{bmatrix} \alpha & 0 \\ 0 & \alpha \end{bmatrix}$ . By (1), then,  $G$  would equal  $\begin{bmatrix} \alpha & 0 \\ 0 & \alpha \end{bmatrix}$  —which it doesn't!

**When is  $\alpha = \beta$ ?** This happens when the discriminant of  $f$  is zero. Its discrim is  $[-\mathcal{S}]^2 - 4 \cdot 1 \cdot \mathcal{P}$ , i.e  $[\alpha^2 + \beta^2 + 2\alpha\beta] - 4\alpha\beta$ . Hence

$$8: \quad \text{Discr}(f) = \mathcal{S}^2 - 4\mathcal{P} \overset{\text{note}}{\equiv} [\alpha - \beta]^2.$$

Since (7) also applies to eigenvalue  $\beta$ , we conclude:

9: Matrix  $G = \begin{bmatrix} \mathcal{S} & -\mathcal{P} \\ 1 & 0 \end{bmatrix}$  is diagonalizable IFF  $G$  has distinct eigenvalues; i.e  $\mathcal{S}^2 \neq 4\mathcal{P}$ .

### Distinct eigenvalues

When  $\alpha \neq \beta$ , our (7) implies that  $G$  is conjugate to diagonal-matrix

$$10: \quad D := \begin{bmatrix} \alpha & 0 \\ 0 & \beta \end{bmatrix} \quad \text{via matrix}$$

$$11: \quad U := \begin{bmatrix} \alpha & \beta \\ 1 & 1 \end{bmatrix}. \quad \text{Note } U^{-1} = \frac{1}{\alpha - \beta} \cdot \begin{bmatrix} 1 & -\beta \\ -1 & \alpha \end{bmatrix}.$$

We populated  $U$  with evecs, using the same order of evals in  $D$ . One can check that  $(U^{-1}GU = D)$ . Hence  $G = UDU^{-1}$ . So for each integer  $n$ ,

$$12: \quad G^n = \frac{1}{\alpha - \beta} \begin{bmatrix} \alpha & \beta \\ 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} \alpha^n & 0 \\ 0 & \beta^n \end{bmatrix} \cdot \begin{bmatrix} 1 & -\beta \\ -1 & \alpha \end{bmatrix}.$$

Multiplying from the right by initial-condition  $\begin{bmatrix} z_1 \\ z_0 \end{bmatrix}$  will produce  $\begin{bmatrix} z_{n+1} \\ z_n \end{bmatrix}$ . Extracting the  $z_n$ , we get the nice formula

$$13: \quad z_n = \frac{1}{\alpha - \beta} \cdot \left[ [z_1 - z_0\beta]\alpha^n - [z_1 - z_0\alpha]\beta^n \right].$$

This formula is symmetric in  $\alpha$  &  $\beta$ , as it must be.

### Equal eigenvalues

Let's take a look at the  $\alpha = \beta$  case. This means that

$$14: \quad G = \begin{bmatrix} 2\alpha & -\alpha^2 \\ 1 & 0 \end{bmatrix}.$$

While we can't conjugate  $G$  to a *diagonal* matrix, we can conjugate to its **Jordan canonical form**. The JCF of (14) is

$$J = J_G := \begin{bmatrix} \alpha & 1 \\ 0 & \alpha \end{bmatrix}.$$

Mysteriously pulling the below  $C$  from a hat, multiplication verifies that

$$15: \quad J = C^{-1}GC, \quad \text{where } C := \begin{bmatrix} \alpha & 1 \\ 1 & 0 \end{bmatrix}$$

and  $C^{-1} = \begin{bmatrix} 0 & 1 \\ 1 & -\alpha \end{bmatrix}$ . Leaving a derivation of  $C$  to later, let's use this to get a formula for  $z_n$ .

Induction on  $n$  shows that

$$J^n := \alpha^{n-1} \cdot \begin{bmatrix} \alpha & n \\ 0 & \alpha \end{bmatrix}, \quad \text{for each } n \in \mathbb{Z}.$$

As before,  $G = CJC^{-1}$  so  $G^n = CJ^nC^{-1}$ . That is,

$$16: \quad \begin{aligned} G^n &= \alpha^{n-1} \cdot \begin{bmatrix} \alpha & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \alpha & n \\ 0 & \alpha \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & -\alpha \end{bmatrix} \\ &= \alpha^{n-1} \cdot \begin{bmatrix} [1+n]\alpha & -n\alpha^2 \\ n & [1-n]\alpha \end{bmatrix}. \end{aligned}$$

Multiplying by  $\begin{bmatrix} z_1 \\ z_0 \end{bmatrix}$ ; the bottom entry in the resulting column-vector is

$$17: \quad z_n = \alpha^{n-1} \cdot \left[ nz_1 + [1-n]\alpha z_0 \right].$$

Filename: Problems/Algebra/LinearAlg/linear-recurr.tex  
As of: Monday 05Mar2007. Typeset: 3Nov2015 at 17:18.