

Liouville's Theorem: Calculus

Jonathan L.F. King

University of Florida, Gainesville FL 32611-2082, USA
squash@ufl.edu

Webpage <http://www.math.ufl.edu/~squash/>

2 February, 2011 (at 21:07)

ABSTRACT: This is Liouville's proof of Liouville's thm on rational approximations of numbers.

Souvenir. The *degree* of an algebraic number α is the degree of the smallest-degree non-zero intpoly having α as a zero.

Whenever I write a rational, e.g. $\frac{p}{q}$, the denominator q is *always positive*.

Warmup. Here is a classic theorem (due probably to Liouville).

1: Theorem. Fix an $\alpha \in \mathbb{R}$. Then

$$\dagger: \quad \left| \alpha - \frac{p}{q} \right| \leq 1/q^2$$

for a seq. of rationals $\frac{p}{q}$ with arbitrarily large q . \diamond

Proof. WLOG α is irrational. Take a large N and look at $0, \alpha, 2\alpha, \dots, [N-1]\alpha$, where we interpret these in the circle group, mod 1. By PHP (Pigeon-hole Principle), for some indices $j < k$ we must have the circle-gp distance $\llbracket k\alpha, j\alpha \rrbracket \leq 1/N$. With $q := k - j$, then, $\llbracket q\alpha, 0 \rrbracket = \llbracket k\alpha, j\alpha \rrbracket \leq 1/N$. I.e., for some integer p ,

$$\ddagger: \quad |q\alpha - p| \leq 1/N \stackrel{\text{note}}{\leq} 1/q.$$

Dividing each side by q yields (\dagger) .

Write the dependency as p_N and q_N . From (\ddagger) , $|\alpha - \frac{p_N}{q_N}| \leq \frac{1}{q_N}$. So $\frac{p_N}{q_N} \rightarrow \alpha$, an irrational, so the set $\{\frac{p_N}{q_N}\}_{N=1}^{\infty}$ is infinite. Thus q_N gets arbitrarily large. \blacklozenge

2: Liouville's Theorem. Suppose α is an irrational but algebraic number. Let $\mathfrak{D} := \text{Deg}(\alpha) \stackrel{\text{note}}{\geq} 2$. Then there exists a posreal C such that for all integers $q > 0$ and p ,

$$3: \quad \left| \alpha - \frac{p}{q} \right| \geq C/q^{\mathfrak{D}}. \quad \blacklozenge$$

Proof. Let $f()$ be a deg- \mathfrak{D} intpoly [necessarily \mathbb{Q} -irreducible] so that $f(\alpha) = 0$. By continuity of f' [the derivative of f] there exists a small interval

$$J := (\alpha - \varepsilon, \alpha + \varepsilon)$$

on which f' is bounded away from infinity. On J then, $1/f'$ is bounded away from zero; by $7/99$, say. Let $C := \text{Min}(\varepsilon, 7/99)$; this is positive.

Consider an arbitrary rational, $\frac{p}{q}$. If $\frac{p}{q} \notin J$ then

$$\left| \alpha - \frac{p}{q} \right| \geq \varepsilon \geq C \geq C/q^{\mathfrak{D}}.$$

So WLOG $\frac{p}{q} \in J$.

Since $f(\frac{p}{q})$ is not zero and f has integer coeffs,

$$4: \quad \left| f\left(\frac{p}{q}\right) \right| \geq 1/q^{\mathfrak{D}}.$$

By the MVT (Mean Value Thm) there is a number ζ between α and $\frac{p}{q}$ such that

$$\left[\alpha - \frac{p}{q} \right] \cdot f'(\zeta) = f(\alpha) - f\left(\frac{p}{q}\right) \stackrel{\text{note}}{=} -f\left(\frac{p}{q}\right).$$

Taking absolute values, then dividing,

$$5: \quad \left| \alpha - \frac{p}{q} \right| = \frac{1}{|f'(\zeta)|} \cdot \left| f\left(\frac{p}{q}\right) \right| \geq \frac{1}{|f'(\zeta)|} / q^{\mathfrak{D}}.$$

But $\frac{p}{q} \in J$, so $\zeta \in J$. Hence (5) implies (3). \blacklozenge

Extension. Liouville's thm holds also when α is rational and $\mathfrak{D} = 1$, as long as we only considers rationals $\frac{p}{q} \neq \alpha$. Simply take J small enough that α is the *only* zero of $f()$ on J ; again (5) holds. \square

Remark. Inequality (3) can be restated as

$$3': \quad \left[\text{Infimum}_{\frac{p}{q} \in \mathbb{Q}} q^{\mathfrak{D}} \cdot \left| \alpha - \frac{p}{q} \right| \right] > 0.$$

This gives a criterion for transcendentality. A number with a sequence such that (6.1) is finite, is called a **Liouville number**. \square

6.0: Transcendental Theorem. Consider a real α . Suppose there exist ∞ ly many rationals p_N/q_N , each **unequal** to α and with $q_N \in [2.. \infty)$, st.

$$6.1: \quad \sup_{N \in \mathbb{Z}_+} [q_N]^N \cdot \left| \alpha - \frac{p_N}{q_N} \right|$$

is finite. Then α is transcendental. \diamond

Proof. Suppose the supremum is 7 . Fixing a posint \mathfrak{D} , could α have degree \mathfrak{D} ? Well, for each $N \geq \mathfrak{D}$ note that

$$[q_N]^{\mathfrak{D}} \cdot \left| \alpha - \frac{p_N}{q_N} \right| \leq 7/[q_N]^{N-\mathfrak{D}} \leq 7/2^{N-\mathfrak{D}}.$$

The RhS goes to zero as $N \nearrow \infty$. This shows the failure of (3'); so α does *not* have degree- \mathfrak{D} . \blacklozenge

7: Example. The following sum of factorial-powers is a Liouville number:

$$\alpha := \sum_{j=1}^{\infty} 1/2^{j!}. \quad \diamond$$

Proof. Fix posint N , let $q := 2^{[N-1]!}$ and define p by $p/q := \sum_{j=1}^{N-1} 1/2^{j!}$. Then $|\alpha - \frac{p}{q}|$ equals

$$\begin{aligned} \sum_{j=N}^{\infty} 1/2^{j!} &= \left[\sum_{j=N}^{\infty} 1/2^{j!-N!} \right] \cdot [1/2^{N!}] \\ &\leq [1 + \frac{1}{2} + \frac{1}{4} + \dots] \cdot [1/2^{N!}] = 2/2^{N!}. \end{aligned}$$

But $q^N = 2^{N!}$, so $q^N \cdot |\alpha - \frac{p}{q}| \leq 2$. Consequently, the number 2 dominates the (6.1) supremum. \blacklozenge

Filename: Problems/Analysis/Calculus/liouville_thm.tex
As of: Thursday 22May2003. Typeset: 2Feb2011 at 21:07.